

LA-UR-19-30720

Approved for public release; distribution is unlimited.

Title: An Effective Field Theory Approach to neutrinoless double beta decay

Author(s): Mereghetti, Emanuele

Intended for: Talk given at the Universita' degli Studi, Milano Bicocca

Issued: 2019-10-21

Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.


An Effective Field Theory approach to neutrinoless $\beta\beta$ decays

Emanuele Mereghetti

with V. Cirigliano, J. Carlson, W. Dekens, J. de Vries, M. Graesser, S. Pastore,
M. Piarulli, U. van Kolck, A. Walker-Loud, R. Wiringa.


arXiv:1710.01729, 1802.10097, 1806.02780, 1907.xxxxx






The Standard Model of Particle Physics


1. describes nature in an economic and elegant way (spontaneously broken) gauge symmetry



$$D_\mu \varphi D^\mu \varphi \rightarrow m_V^2 V_\mu V^\mu$$



$$\bar{q}_L \varphi d_R \rightarrow m_d \bar{d}_L d_R$$




The Standard Model of Particle Physics

- generates fermion and boson masses via the Higgs mechanism

confirmed at the LHC!¹²

¹... for gauge bosons and heaviest fermions ...

²... SM flavor structure is somewhat mysterious ...



neutrino oscillation experiments:

- neutrino have masses too!
- well described by 2 mass differences


& 3×3 unitary mixing matrix U_{PMNS}

$\Delta m_{ij}^2, |U_{li}|$ well known

CP (and Majorana) phases, mass ordering ?

Introduction

$$|U_{PMNS}| = \begin{pmatrix} \nu_1 & \nu_2 & \nu_3 \\ \nu_e & \text{blue square} & \text{red square} & \text{small pink square} \\ \nu_\mu & \text{green square} & \text{red square} & \text{orange square} \\ \nu_\tau & \text{green square} & \text{red square} & \text{orange square} \end{pmatrix}$$



neutrino oscillation experiments:


- neutrino have masses too!
- well described by 2 mass differences

& 3×3 unitary mixing matrix U_{PMNS}

$\Delta m_{ij}^2, |U_{ii}|$ well known

CP (and Majorana) phases, mass ordering ?

Introduction



$$\begin{array}{c} \longrightarrow \\ \nu_L \quad \bullet \quad \nu_R \\ \longrightarrow \end{array}$$

Dirac

$$m_i \bar{\nu}_R^i \nu_L^i$$


$$\begin{array}{c} \longrightarrow \\ \nu_L \quad \blacksquare \quad \nu_L \\ \longleftarrow \end{array}$$

Majorana

$$m_i \nu_L^{T i} C \nu_L^i$$

- ... but neutrinos are peculiar ...
- two possible mass terms

Introduction



Dirac



Majorana

- ... but neutrinos are peculiar ...
- two possible mass terms

Dirac:

- no ν_R in the SM
- if ν_R what forbids

$$m_{\nu_R} \nu_R^T C \nu_R ?$$

new mixings in oscillation exp.



Majorana:

- break $SU(2)_L$
- restore $SU(2)_L$

$$\frac{1}{\Lambda} \varepsilon_{ij} \varepsilon_{mn} L_i^T C L_m H_j H_n$$

at the price of heavy new physics

Introduction




M. Duerr, M. Lindner, K. Zuber, '11

- double beta decay is rare doubly-weak decay process
- $2\nu\beta\beta$ allowed in SM


$$T_{1/2}^{2\nu}(^{76}\text{Ge} \rightarrow ^{76}\text{Se}) = (1.84_{-0.10}^{+0.14}) \times 10^{21} \text{ yr}$$

GERDA coll., '15

Introduction



M. Duerr, M. Lindner, K. Zuber, '11



- $0\nu\beta\beta$ violates lepton number L by two units

$$(A, Z) \rightarrow (A, Z + 2) + e^- e^-$$


possible iff ν s have a Majorana mass

- L conserved in the SM, $0\nu\beta\beta$


new physics!

implications for ν mass, leptogenesis ...

Introduction



- current bounds already impressive
- next-generation tonne-scale experiments will improve by 1-2 orders




Next generation of experiments sensitive to a variety of LNV scenarios

1. LNV originates at very high scales


direct connection between ν oscillations and $0\nu\beta\beta$

Introduction



Next generation of experiments sensitive to a variety of LNV scenarios

1. LNV originates at very high scales
 - direct connection between ν oscillations and $0\nu\beta\beta$
 - clear interpretative framework and goals
 - e.g. rule out inverted hierarchy




Next generation of experiments sensitive to a variety of LNV scenarios

2. LNV at intermediate scales

$0\nu\beta\beta$ is mediated by new particles
could be accessible at colliders

general framework to interpret $0\nu\beta\beta$ exp.?
with controlled uncertainties ?

EFT approach to LNV



EFT approach to LNV

- half-life anatomy

$$\left(T_{1/2}^{0\nu}\right)^{-1} = \frac{m_{\beta\beta}^2}{m_e^2} G_{01} g_A^4 |M^{0\nu}|^2 + \dots \quad M^{0\nu} = \langle 0^+ | V_\nu | 0^+ \rangle$$

- fundamental LNV parameter
- “phase space factors”
leptonic structure of LNV operators
- from quarks to hadrons
- nuclear matrix elements of “ ν potentials”

What EFTs can do:


parametrize $0\nu\beta\beta$ w. few coefficients
that can be matched to models

identify QCD input & its uncertainty

systematically derive the ν potentials

check NMEs in simpler systems

EFT approach to LNV



J. Engel and J. Menéndez, '16


Final goal:

- help reduce theory uncertainties on $M^{0\nu}$

... not quite there yet ...

The Standard Model as an EFT

The Standard Model as an Effective Field Theory



Write down all possible operators with

- SM fields
- local $SU(3)_c \times SU(2)_L \times U(1)_Y$ invariance
- dimension ≤ 4

$m_\nu = 0$
no ΔL interactions

assume no light sterile ν_R

The Standard Model as an EFT

- why stop at dim=4?

$$\mathcal{L} = \mathcal{L}_{SM} + \sum \frac{c_{i,5}}{\Lambda} \mathcal{O}_{5i} + \sum \frac{c_{i,6}}{\Lambda^2} \mathcal{O}_{6i} + \sum \frac{c_{i,7}}{\Lambda^3} \mathcal{O}_{7i} + \dots$$

$\Lambda \gg v = 246 \text{ GeV}$

- \mathcal{O} are gauge invariant
- no need to impose accidental symmetries as L

The Standard Model as an EFT


- why stop at dim=4?

$$\mathcal{L} = \mathcal{L}_{SM} + \sum \frac{c_{i,5}}{\Lambda} \mathcal{O}_{5i} + \sum \frac{c_{i,6}}{\Lambda^2} \mathcal{O}_{6i} + \sum \frac{c_{i,7}}{\Lambda^3} \mathcal{O}_{7i} + \dots$$

$\Lambda \gg v = 246 \text{ GeV}$

- \mathcal{O} are gauge invariant
- no need to impose accidental symmetries as L
- **one** dimension 5 operator

S. Weinberg, '79







$$\frac{1}{\Lambda} \varepsilon_{ij} \varepsilon_{mn} L_i^T C L_m H_j H_n \rightarrow \frac{v^2}{\Lambda} \nu_L^T C \nu_L$$


neutrino masses and mixings

$$\Lambda \sim 10^{14} \text{ GeV}$$

LNV at dim. 7, dim. 9



$$\left(\frac{\nu}{\Lambda}\right)^3$$









$$\left(\frac{\nu}{\Lambda}\right)^5$$

- dim.7 operators mostly induce β decay with “wrong” ν

\implies long range contribs. to $0\nu\beta\beta$

- dim. 9 induce short-range contributions to $0\nu\beta\beta$



Connection to models



- specific models will match onto one or several operators
- e.g. LR symmetric model
dim. 5, 7 & 9 (with different Yukawas)


can match any model to EFT

Dimension 7 at LHC



- some limits from $pp \rightarrow e\nu$: $\Lambda \lesssim 2.5 \text{ TeV}$
- no way to disentangle from $\Delta L = 0$ non-standard couplings
- no way to tell Dirac from Majorana

Dimension 7 at LHC



- to be sure is $\Delta L = 2$:
analyze the neutrino with another weak interaction

$$pp \rightarrow e^- e^- W^+ (e^+ \nu)$$


$$pp \rightarrow e^- e^- W^+ (jj)$$

$$pp \rightarrow e^- e^- 2j$$


- non competitive for dim. 7 operators
- study for dim. 9

Low-energy EFT for LNV


$\Delta L = 2$ Lagrangian at 1 GeV



$$\frac{v}{\Lambda}$$



$$\frac{v^3}{\Lambda^3}$$



$$\frac{v^3}{\Lambda^3}, \frac{v^5}{\Lambda^5}$$


$$\mathcal{L}_{\Delta L=2}(\nu, e, u, d) = -\frac{1}{2}(m_\nu)_{ij}\nu^T C\nu^j + C_\Gamma \nu^T C \Gamma e \mathcal{O}_\Gamma + C_{\Gamma'} e^T C \Gamma' e \mathcal{Q}_{\Gamma'}$$

quark bilinear

four-quark

- match onto a EFT for nucleons?

Interlude: Chiral Effective Field Theory



$$\mathcal{L} = -\frac{1}{4} G_{\mu\nu}^a G^{a\mu\nu} + \bar{q}_L i \not{D} q_L + \bar{q}_R i \not{D} q_R - \bar{q}_L \mathcal{M} q_R - \bar{q}_R \mathcal{M} q_L + \mathcal{L}_{\Delta L=2} + \dots$$

Chiral symmetry & its spontaneous breaking:

- pions are pseudo-Goldstone
 - a. light, $m_\pi \ll \Lambda_\chi = 4\pi F_\pi \sim 1 \text{ GeV}$
 - b. and weakly coupled
- EFT expansion in powers of $\{Q, m_\pi\}/\Lambda_\chi$

$$\frac{dm_N}{dm_\pi^2} =$$

~ 1

$\sim \frac{m_\pi}{4\pi F_\pi}$

$\sim \frac{m_\pi^2}{(4\pi F_\pi)^2}$

- only one relevant scale $Q \sim m_\pi$
 “HQET”-like w. “soft” pion & nucleon modes $p \sim (m_\pi, m_\pi)$
- perturbative expansion of χ EFT Lagrangian


$$\mathcal{L} = \mathcal{L}^{(0)} + \frac{1}{\Lambda_\chi} \mathcal{L}^{(1)} + \frac{1}{(\Lambda_\chi)^2} \mathcal{L}^{(2)} \dots$$

- and amplitudes

$$\mathcal{A} = \sum a_n \left(\frac{Q}{\Lambda_\chi} \right)^n$$


derivatives, pion masses & loops suppressed by Q/Λ_χ

Chiral EFT. $A \geq 2$



1. another scale in the problem: Q^2/m_N
 “NRQCD”-like with “potential” $p \sim (Q^2/m_N, Q)$ modes

- resum an infinite class of diagrams
- equivalent to solving Lippmann-Schwinger equation



1. another scale in the problem: Q^2/m_N


“NRQCD”-like with “potential” $p \sim (Q^2/m_N, Q)$ modes

- resum an infinite class of diagrams
- equivalent to solving Lippmann-Schwinger equation



- another scale in the problem: Q^2/m_N
 “NRQCD”-like with “potential” $p \sim (Q^2/m_N, Q)$ modes
 - resum an infinite class of diagrams
 - equivalent to solving Lippmann-Schwinger equation
- V is organized in powers of $\frac{Q}{\Lambda_\chi}$
- the LO potential is singular (“short-range core”)
 - new divergences in solution of LS
 - can invalidate power counting based on NDA

Nuclear EFT(s)



from D. R. Entem and R. Machleidt, '17

see also:

P. Reinert, H. Krebs, E. Epelbaum, '18

M. Piarulli *et al.*, '16


M. Piarulli *et al.*, '14

A. Nogga, R. Timmermans, B. van Kolck, '05

D. Kaplan, M. Savage, M. Wise, '96

- LECs are fit to data in 2- and 3-nucleon systems
- and predict light-nuclear observables


Nuclear EFT(s)



- LECs are fit to data in 2- and 3-nucleon systems
- and predict light-nuclear observables


WARNING: unresolved issues with power counting

External currents in chiral EFT



- similar expansions for external currents
- consistent interactions & currents:
ab initio solution to long-standing problem in β decays
 “ g_A quenching”

External currents in chiral EFT




P. Gysbers *et al.*, '19

- similar expansions for external currents
- consistent interactions & currents:
ab initio solution to long-standing problem in β decays
“ g_A quenching”

Revisiting the light Majorana- ν exchange mechanism

Chiral EFT approach to light- ν exchange mechanism



$$\mathcal{L} = \mathcal{L}_{\text{QCD}} - \frac{4G_F}{\sqrt{2}} \bar{u}_L \gamma^\mu d_L \bar{e}_L \gamma_\nu \nu_L - \frac{m_{\beta\beta}}{2} \nu_L^T C \nu_L$$


- weak currents are mainly one-body
- $0\nu\beta\beta$ mediated by exchange of “potential” neutrinos

$$V_\nu = \mathcal{A} \tau^{(1)+} \tau^{(2)+} + \frac{1}{\mathbf{q}^2} \left\{ \mathbf{1}^{(a)} \times \mathbf{1}^{(b)} - g_A^2 \boldsymbol{\sigma}^{(a)} \cdot \boldsymbol{\sigma}^{(b)} \left(\frac{2}{3} + \frac{1}{3} \frac{m_\pi^4}{(\mathbf{q}^2 + m_\pi^2)^2} \right) + \dots \right\}.$$

$$\mathcal{A} = 2G_F^2 m_{\beta\beta} \bar{e}_L C \bar{e}_L^T$$

agrees with all $0\nu\beta\beta$ literature

Standard mechanism. Higher orders



At N²LO $\mathcal{O}(\mathbf{q}^2/\Lambda_\chi^2)$


1. correction to the one-body currents (magnetic moment, radii, ...)

$$g_A(\mathbf{q}^2) = g_A \left(1 - r_A^2 \frac{\mathbf{q}^2}{6} + \dots \right) \quad r_A = 0.47(7) \text{ fm}$$

2. two-body corrections to V and A currents
3. pion-neutrino loops & local counterterms

UV divergences signal short-range sensitivity at N²LO

Standard mechanism. Higher orders



At $N^2\text{LO}$ $\mathcal{O}(\mathbf{q}^2/\Lambda_\chi^2)$

1. correction to the one-body currents (magnetic moment, radii, ...)


$$g_A(\mathbf{q}^2) = g_A \left(1 - r_A^2 \frac{\mathbf{q}^2}{6} + \dots \right) \quad r_A = 0.47(7)\text{fm}$$

2. two-body corrections to V and A currents
3. pion-neutrino loops & local counterterms

UV divergences signal short

WARNING: based on naive dimensional analysis
“Weinberg’s counting”

Is the Weinberg counting consistent for $0\nu\beta\beta$?



- Weinberg's counting fails in 1S_0 channel

D. Kaplan, M. Savage, M. Wise, '96


- study $nn \rightarrow ppe^-e^-$ with LO χ EFT strong potential

$$V_{\text{strong}}(r) = \tilde{C} \delta^{(3)}(\mathbf{r}) + \frac{g_A^2 m_\pi^2}{16\pi F_\pi^2} \frac{e^{-m_\pi r}}{4\pi r}$$

\tilde{C} fit to 1S_0 scattering length

- no problem with Yukawa potential

Is the Weinberg counting consistent for $0\nu\beta\beta$?



- Weinberg's counting fails in 1S_0 channel

D. Kaplan, M. Savage, M. Wise, '96


- study $nn \rightarrow ppe^-e^-$ with LO χ EFT strong potential

$$V_{\text{strong}}(r) = \tilde{C} \delta^{(3)}(\mathbf{r}) + \frac{g_A^2 m_\pi^2}{16\pi F_\pi^2} \frac{e^{-m_\pi r}}{4\pi r}$$

\tilde{C} fit to 1S_0 scattering length

- no problem with Yukawa potential
- and one insertion of short-range potential

Inconsistency of the Weinberg counting




$$\frac{1}{2}(1 + 2g_A^2) \left(\frac{m_N \tilde{C}}{4\pi} \right)^2 \left(\frac{1}{\epsilon} + \log \mu^2 \right)$$

- two-loop diagrams w. two insertions of \tilde{C} have UV log divergence

need a local LNV counterterm at LO!

Inconsistency of the Weinberg counting



- two-loop diagrams w. two insertions of \tilde{C} have UV log divergence

need a local LNV counterterm at LO!


- renormalization requires to modify the LO ν potential

$$V_{\text{LNV}} = V_\nu - 2g_\nu \tau^{(1)+} \tau^{(2)+} \mathcal{A}$$

- the coupling g_ν is larger than NDA

$$g_\nu \sim \frac{1}{F_\pi^2} \gg \frac{1}{(4\pi F_\pi)^2}$$

Inconsistency of the Weinberg counting



$$\mathcal{A}_\nu = \int d^3\mathbf{r} \psi_{\mathbf{p}'}^-(\mathbf{r}) V_\nu(\mathbf{r}) \psi_{\mathbf{p}}^+(\mathbf{r})$$

$$g_\nu = \left(\frac{m_N \tilde{C}}{4\pi} \right)^2 \tilde{g}_\nu$$


$$\tilde{g}_\nu \sim b - \frac{1}{2}(1 + 2g_A^2) \log R_S$$

- divergence is not an artifact of dim. reg.
e.g use a gaussian cut-off


$$\tilde{C} \delta^{(3)}(\mathbf{r}) \rightarrow \frac{\tilde{C}}{\pi^{3/2} R_S^3} e^{-r^2/R_S^2}$$

- \mathcal{A}_ν shows logarithmic dependence on R_S (+ power corrections)


Relation between $0\nu\beta\beta$ and EM isospin breaking




- the dynamics of QCD seems to imply a short-range component for V_ν
 - does this happen anywhere else? How to fix finite piece of the coupling?
- Charge independence breaking in NN scattering




Relation between $0\nu\beta\beta$ and EM isospin breaking



- the dynamics of QCD seems to imply a short-range component for V_ν
 - does this happen anywhere else? How to fix finite piece of the coupling?
- Charge independence breaking in NN scattering



Relation between $0\nu\beta\beta$ and EM isospin breaking




- only two $I = 2$ operators w. same properties as weak/EM currents

$$\begin{aligned} \mathcal{L}_{I=2} &= c C_1 \left(\bar{N} Q_L N \bar{N} Q_L N - \frac{\text{Tr}[Q_L^2]}{6} \bar{N} \boldsymbol{\tau} N \cdot \bar{N} \boldsymbol{\tau} N + L \rightarrow R \right) \\ &+ c C_2 \left(\bar{N} Q_L N \bar{N} Q_R N - \frac{\text{Tr}[Q_L Q_R]}{6} \bar{N} \boldsymbol{\tau} N \cdot \bar{N} \boldsymbol{\tau} N + L \rightarrow R \right) \\ Q_L &= u^\dagger Q_L u \quad Q_R = u Q_R u^\dagger, \quad u = 1 + \frac{i\boldsymbol{\pi} \cdot \boldsymbol{\tau}}{2F_\pi} + \dots \end{aligned}$$

- weak interactions:** $Q_L = \tau^+$, $Q_R = 0$, $c_{LNV} = 2G_F^2 m_{\beta\beta} \bar{e}_L C \bar{e}_L^T$
- EM interactions:** $Q_L = \frac{\tau^z}{2}$, $Q_R = \frac{\tau^z}{2}$, $c_{e^2} = \frac{e^2}{4}$

Relation between $0\nu\beta\beta$ and EM isospin breaking




- only two $I = 2$ operators w. same properties as weak/EM currents

$$\begin{aligned}
 \mathcal{L}_{I=2} &= c C_1 \left(\bar{N} Q_L N \bar{N} Q_L N - \frac{\text{Tr}[Q_L^2]}{6} \bar{N} \boldsymbol{\tau} N \cdot \bar{N} \boldsymbol{\tau} N + L \rightarrow R \right) \\
 &+ c C_2 \left(\bar{N} Q_L N \bar{N} Q_R N - \frac{\text{Tr}[Q_L Q_R]}{6} \bar{N} \boldsymbol{\tau} N \cdot \bar{N} \boldsymbol{\tau} N + L \rightarrow R \right) \\
 Q_L &= u^\dagger Q_L u \quad Q_R = u Q_R u^\dagger, \quad u = 1 + \frac{i\boldsymbol{\pi} \cdot \boldsymbol{\tau}}{2F_\pi} + \dots
 \end{aligned}$$

- $C_1 = g_\nu$ by chiral symmetry!
- C_1 and C_2 differ at multiplication level

cannot disentangle in NN scattering
but give an idea of $0\nu\beta\beta$ counterterm

Weinberg counting for isospin breaking operators



- leading $I = 2$ potential in 1S_0 channel from γ exchange & pion mass splitting


$$V_{\text{CIB}} = \frac{e^2}{4} \left(\tau_3^{(1)} \tau_3^{(2)} - \frac{1}{3} \boldsymbol{\tau}^{(1)} \cdot \boldsymbol{\tau}^{(2)} \right) \frac{1}{\mathbf{q}^2} \left\{ 1 - \frac{g_A^2}{F_\pi^2} \frac{\Delta m_\pi^2}{e^2} \boldsymbol{\sigma}^{(1)} \cdot \mathbf{q} \boldsymbol{\sigma}^{(2)} \cdot \mathbf{q} \frac{\mathbf{q}^2}{(\mathbf{q}^2 + m_\pi^2)^2} \right\}.$$

$$\Delta m_\pi^2 = m_{\pi^\pm}^2 - m_{\pi^0}^2$$

- short-range contributions suppressed

$$V_{\text{CIB}}^S = \frac{e^2}{2} \frac{C_1 + C_2}{2} \left(\tau^{(1)z} \tau^{(2)z} - \frac{1}{3} \boldsymbol{\tau}^{(1)} \cdot \boldsymbol{\tau}^{(2)} \right) \quad C_1 \sim C_2 \sim \frac{1}{(4\pi F_\pi)^2}$$


Relation to charge-independence breaking



- fit one charge-independent \tilde{C} in np
- compute a_{nn} and a_C

log divergence! need a ct in each channel

Relation to charge-independence breaking




1. LO analysis of isospin breaking show log dependence

$$\frac{C_1 + C_2}{2} = \left(\frac{m_N \tilde{C}}{4\pi} \right)^2 \frac{\tilde{C}_1 + \tilde{C}_2}{2} \sim_{R_S=0.5} \frac{16}{(4\pi F_\pi)^2}$$

disagree with Weinberg's counting!

2. all high-quality chiral & pheno NN potentials include short-range CIB

Impact on $0\nu\beta\beta$ nuclear matrix elements




$nn \rightarrow pp$

- assume $C_1(R_S) = C_2(R_S)$
- LNV matrix element is scale independent
- effect of short-range potential $\sim 10\%$

$\Delta I = 0$ transition

Impact on $0\nu\beta\beta$ nuclear matrix elements




- extract CIB potential V_{CIB}^S from AV18 or χEFT (rescaled by c_{LNV}/c_{e^2})

$$\text{AV18: } M_L = 7.45, \quad M_S = 0.48$$

$$\chi\text{EFT: } M_L = 7.82, \quad M_S = 1.15$$

$\sim 10\%$ corrections

Impact on $0\nu\beta\beta$ nuclear matrix elements



- larger corrections to $I = 2$ transitions



$$\text{AV18: } M_L = 0.653, \quad M_S = 0.518$$

$$\chi\text{EFT: } M_L = 0.725, \quad M_S = 0.533$$

> 50% corrections

- ... but uncontrolled theory error from assuming $C_1 = C_2!$

Impact on $0\nu\beta\beta$ nuclear matrix elements



- larger corrections to $I = 2$ transitions


$$\text{AV18: } M_L = 0.653, \quad M_S = 0.518$$

$$\chi\text{EFT: } M_L = 0.725, \quad M_S = 0.533$$

> 50% corrections

- ... but uncontrolled theory error from assuming $C_1 = C_2!$

Partial summary: light Majorana exchange



- chiral EFT uncovers additional source of uncertainty
QCD at $\Lambda \sim 1$ GeV gives non-negligible contributions to V_ν
confirmed by isospin breaking in NN scattering
- V_ν needs non-ptb matching between QCD & chiral EFT
e.g. Lattice QCD calculation of $nn \rightarrow ppe^-e^-$


first results on $\pi^- \rightarrow \pi^+ e^- e^-$

Xu Feng, *et al.* '18


- problem only in the 1S_0 channel
Weinberg's counting ok in $^3P_J, ^1D_2$ channels

Chiral EFT for non-standard mechanisms

Chiral EFT for non-standard mechanisms



Dim. 9 operators






1. LL LL : $\mathcal{O}_1 = \bar{u}_L \gamma^\mu d_L \bar{u}_L \gamma_\mu d_L$
2. LR LR : $\mathcal{O}_2 = \bar{u}_L d_R \bar{u}_L d_R$, $\mathcal{O}_3 = \bar{u}_L^\alpha d_R^\beta \bar{u}_L^\beta d_R^\alpha$
3. LL RR : $\mathcal{O}_4 = \bar{u}_L \gamma^\mu d_L \bar{u}_R \gamma_\mu d_R$, $\mathcal{O}_5 = \bar{u}_L^\alpha \gamma^\mu d_L^\beta \bar{u}_R^\beta \gamma_\mu d_R^\alpha$

- several unjustified assumptions in the literature . . .

$$\text{e.g. } \langle pp | \bar{u}_L \gamma^\mu d_L \bar{u}_R \gamma_\mu d_R | nn \rangle = \langle p | \bar{u}_L \gamma^\mu d_L | n \rangle \langle p | \bar{u}_R \gamma_\mu d_R | n \rangle = (1 - 3g_A^2)$$

inconsistent with QCD, miss chiral dynamics

LNV interactions from dim. 9 operators



- πN couplings, only important for \mathcal{O}_1
- NN couplings


$$\mathcal{L}_{NN} = \left(g_1^{NN} C_{1L}^{(9)} + g_2^{NN} C_{2L}^{(9)} + g_3^{NN} C_{3L}^{(9)} + g_4^{NN} C_{4L}^{(9)} + g_5^{NN} C_{5L}^{(9)} \right) (\bar{p}n) (\bar{p}n) \frac{\bar{e}_L C \bar{e}_L^T}{\nu^5}$$

- size depends on chiral properties of $\mathcal{O}_{1,\dots,5}$

$$g_1^{NN} \sim \mathcal{O}(1), \quad g_{2,3,4,5}^{NN} \sim \mathcal{O}\left(\frac{\Lambda_\chi^2}{F_\pi^2}\right)$$

enhanced w.r.t NDA!

$\pi\pi$ matrix elements



$$g_1^{\pi\pi} = +0.4$$

$$g_2^{\pi\pi} = -(1.8 \text{ GeV})^2$$

$$g_3^{\pi\pi} = +(1.0 \text{ GeV})^2$$

$$g_4^{\pi\pi} = -(1.7 \text{ GeV})^2$$

$$g_5^{\pi\pi} = -(3.6 \text{ GeV})^2$$


A. Nicholson *et al.*, CalLat collaboration, '18

- $\pi\pi$ matrix elements well determined in LQCD

good agreement with NDA

- $nn \rightarrow pp$ will allow to determine g_i^{NN} and test the chiral EFT power counting

$0\nu\beta\beta$ potential



- NME differ dramatically from factorization
e.g $C_4^{(9)}$

$$M = -\frac{g_4^{\pi\pi} C_4^{(9)}}{2m_N^2} \left(\frac{1}{2} M_{AP,sd}^{GT} + M_{PP,sd}^{GT} \right) \sim -0.60 C_4^{(9)}$$

$$M_{\text{fact}} = -\frac{3g_A^2 - 1}{2g_A^2} \frac{m_\pi^2}{m_N^2} C_4^{(9)} M_{F,sd} \sim -0.04 C_4^{(9)}$$

bigger error than from NMEs ...

Master Formula

$$\begin{aligned} \left(T_{1/2}^{0\nu}\right)^{-1} &= g_A^4 \left\{ G_{01} \left(|\mathcal{A}_\nu|^2 + |\mathcal{A}_R|^2 \right) - 2(G_{01} - G_{04}) \operatorname{Re} \mathcal{A}_\nu^* \mathcal{A}_R + 4G_{02} |\mathcal{A}_E|^2 \right. \\ &\quad + 2G_{04} \left[|\mathcal{A}_{m_e}|^2 + \operatorname{Re} \left(\mathcal{A}_{m_e}^* (\mathcal{A}_\nu + \mathcal{A}_R) \right) \right] - 2G_{03} \operatorname{Re} \left[(\mathcal{A}_\nu + \mathcal{A}_R) \mathcal{A}_E^* + 2\mathcal{A}_{m_e} \mathcal{A}_E^* \right] \\ &\quad \left. + G_{09} |\mathcal{A}_M|^2 + G_{06} \operatorname{Re} \left[(\mathcal{A}_\nu - \mathcal{A}_R) \mathcal{A}_M^* \right] \right\}. \end{aligned}$$


- G_{01}, \dots, G_{09} : phase space factors
leptonic structure of the operators
- $\mathcal{A}_\nu, \dots, \mathcal{A}_M$: combinations of NME and couplings

$$\mathcal{A}_\nu = \frac{m_{\beta\beta}}{m_e} \mathcal{M}_\nu^{(3)} + \frac{m_N}{m_e} \mathcal{M}_\nu^{(6)} + \frac{m_N^2}{m_{e\nu}} \mathcal{M}_\nu^{(9)}$$

- organized in powers of Λ_χ/Λ and Q/Λ_χ

$$\mathcal{M}_\nu^{(9)} = -\frac{1}{2m_N^2} C_{\pi\pi L}^{(9)} \left(\frac{1}{2} M_{GT, sd}^{AP} + M_{GT, sd}^{PP} + \frac{1}{2} M_{T, sd}^{AP} + M_{T, sd}^{PP} \right) + \dots$$

Nuclear matrix elements



calculations differ by
factor of 2-3


- at LO in χ EFT, **all** nuclear matrix elements (NME) can be expressed in terms of existing calculations
- 8 long-range NME
- 6 short-range NME

contribute to light ν exchange

contribute to heavy Majorana ν exchange


Phenomenology

Phenomenology



- $0\nu\beta\beta$ put strong limits on dim. 7 operators
- dim. 9 in the TeV range


pattern can be understood from effective dimension
& chiral properties of $0\nu\beta\beta$ operator



- $0\nu\beta\beta$ put strong limits on dim. 7 operators
- dim. 9 in the TeV range

pattern can be understood from effective dimension
& chiral properties of $0\nu\beta\beta$ operator

$0\nu\beta\beta$ in the Left-Right Symmetric Model





- generate dim. 5, 7 and 9
- dim. 7 and dim. 9 are chirally suppressed

Case 1 $m_{W_R} = 4.5 \text{ TeV}$, $m_{\Delta_R} = 10 \text{ TeV}$, $U_R = U_{\text{PMNS}}$,

$$m_{\nu_R} \sim m_{W_R}$$

- strong collider bounds on m_{W_R} suppress dim. 7 and dim. 9 contris.
- light- ν Majorana mass dominates in IH
- dim. 9 sizable in NH, but not in reach

$0\nu\beta\beta$ in the LRSM




Case 2 $m_{W_R} = 4.5$ TeV, $m_{\Delta_R} = 10$ TeV, $U_R = U_{\text{PMNS}}$,

$$m_{\nu_R} \sim 10 \text{ GeV}$$

- not ruled out by LEP, LHC searches
- dim. 9 contribution becomes dominant
- in conflict with current $0\nu\beta\beta$ limits

$0\nu\beta\beta$ in the LRSM



- disentangle LRSM from standard mechanism?
- different isotopes are largely degenerate
- electron energy and angular distributions as well
- need interplay with LHC searches!

Summary

(χ)EFTs & $0\nu\beta\beta$ decay:

- connection with collider observables
- model-independent parameterization of low-energy $\Delta L = 2$ operators
- systematic organization of ν potentials

Standard mechanism:

- LO short-range potential missing from all existing calculations

additional $\mathcal{O}(1)$ contribution


- g_ν can be guessed from isospin breaking in NN scattering
new LQCD calculations required

Non Standard mechanisms:

- identify QCD input and hadronic uncertainties
- derive minimal sets on neutrino potentials & relate to existing calculations

Backup

Usoft contribution to the amplitude



overlap $\langle n|J_\mu|i\rangle$
same as in $2\nu\beta\beta!$

4. soft neutrinos, which couple to the nuclear bound states

$$T_{\text{usoft}}(\mu_{\text{us}}) = \frac{T_{\text{lept}}}{8\pi^2} \sum_n \langle f|J_\mu|n\rangle \langle n|J^\mu|i\rangle \left\{ (E_2 + E_n - E_i) \left(\log \frac{\mu_{\text{us}}}{2(E_2 + E_n - E_i)} + 1 \right) + 1 \leftrightarrow 2 \right\}$$

- corrections to the “closure approximation”
- suppressed by $E/(4\pi k_F)$

Is the Weinberg counting consistent?

$$\begin{aligned}
 iA &= \text{[diagram: tree-level exchange]} + \text{[diagram: one-loop exchange]} + \text{[diagram: two-loop exchange]} + \dots \\
 &= \text{[diagram: tree-level exchange]} + \frac{\text{[diagram: one-loop exchange]} + \text{[diagram: two-loop exchange]}}{1 - \text{[diagram: one-loop exchange]}}
 \end{aligned}$$

$m_\pi^2 \left(\frac{1}{\epsilon} + \log \mu^2 \right)$

D. Kaplan, M. Savage, M. Wise, '96

- NDA does not work in NN scattering
- m_π dependence of short-range nuclear force should be subleading

$$\mathcal{L} = -\tilde{C}(N^T P^1 S_0 N)(N^T P^1 S_0 N)^\dagger - \frac{m_\pi^2}{(4\pi F_\pi)^2} D_2(N^T P^1 S_0 N)(N^T P^1 S_0 N)^\dagger + \dots$$

$$4\pi F_\pi = \Lambda_\chi \sim 1 \text{ GeV}$$

Is the Weinberg counting consistent?

$$\begin{aligned}
 iA &= \text{[diagram: tree-level exchange]} + \text{[diagram: one-loop exchange]} + \text{[diagram: two-loop exchange]} + \dots \\
 &= \text{[diagram: tree-level exchange]} + \frac{\text{[diagram: one-loop exchange]} + \text{[diagram: two-loop exchange]}}{1 - \text{[diagram: one-loop exchange]}}
 \end{aligned}$$

$m_\pi^2 \left(\frac{1}{\epsilon} + \log \mu^2 \right)$

D. Kaplan, M. Savage, M. Wise, '96

- NDA does not work in NN scattering
- m_π dependence of short-range nuclear force should be subleading

$$\mathcal{L} = -\tilde{C}(N^T P^1 S_0 N)(N^T P^1 S_0 N)^\dagger - \frac{m_\pi^2}{(4\pi F_\pi)^2} D_2 (N^T P^1 S_0 N)(N^T P^1 S_0 N)^\dagger + \dots$$


$$4\pi F_\pi = \Lambda_\chi \sim 1 \text{ GeV}$$

- ... but UV divergences in the LO amplitude require a promotion ...

conflict between NDA & short-range core of nuclear force

Low-energy Effective Lagrangian for $\Delta L = 2$

$\Delta L = 2$ Lagrangian at 1 GeV




$$\mathcal{L}_{\Delta L=2} = \mathcal{L}_{\Delta L=2}^{\Delta e=0} + \mathcal{L}_{\Delta L=2}^{\Delta e=1} + \mathcal{L}_{\Delta L=2}^{\Delta e=2}$$

- $\mathcal{L}_{\Delta L=2}^{\Delta e=0}$ includes ν masses, magnetic moments, ...

$$\mathcal{L}_{\Delta L=2} = -\frac{1}{2}(m_\nu)_{ij} \nu_L^{Tj} C \nu_L^i + \dots \quad m_\nu \sim \mathcal{O}\left(\frac{v^2}{\Lambda}\right)$$

$\Delta L = 2$ Lagrangian at 1 GeV




$$\mathcal{L}_{\Delta L=2} = \mathcal{L}_{\Delta L=2}^{\Delta e=0} + \mathcal{L}_{\Delta L=2}^{\Delta e=1} + \mathcal{L}_{\Delta L=2}^{\Delta e=2}$$

- $\mathcal{L}_{\Delta L=2}^{\Delta e=1}$ starts at dim. 6, $C_i^{(6)} = \mathcal{O}\left(\frac{v^3}{\Lambda^3}\right)$

$$\mathcal{L}_{\Delta L=2}^{(6)} = \frac{2G_F}{\sqrt{2}} \left\{ C_{\text{VL}}^{(6)} \bar{d}_L \gamma^\mu u_L \nu_L^T C \gamma_\mu e_R + C_{\text{VR}}^{(6)} \bar{d}_R \gamma^\mu u_R \nu_L^T C \gamma_\mu e_R \right. \\ \left. + C_{\text{SL}}^{(6)} \bar{d}_R u_L \nu_L^T C e_L + C_{\text{SR}}^{(6)} \bar{d}_L u_R \nu_L^T C e_L + C_{\text{T}}^{(6)} \bar{d}_R \sigma^{\mu\nu} u_L \nu_L^T C \sigma_{\mu\nu} e_L \right\}$$

β decay w. the “wrong” neutrino & all possible Lorentz structures

$\Delta L = 2$ Lagrangian at 1 GeV



- $\mathcal{L}_{\Delta L=2}^{\Delta e=2}$ starts at dim. 9


$$\mathcal{L}_{\Delta L=2}^{(9)} = \frac{2G_F^2}{v} \left[\sum_{i=\text{scalar}} \left(C_i^{(9)} \bar{e}_L C \bar{e}_L^T + C_i^{(9)'} \bar{e}_R C \bar{e}_R^T \right) O_i + \bar{e}_R \gamma_\mu C \bar{e}_L^T \sum_{i=\text{vector}} C_{iV}^{(9)} O_i^\mu \right]$$

- a small set receives contributions from dim. 7 operators

$$C_1^{(9)}, C_{4,5}^{(9)} \sim \mathcal{O} \left(\frac{v^3}{\Lambda^3} \right), \quad C_i^{(9)'} \sim \mathcal{O} \left(\frac{v^5}{\Lambda^5} \right)$$

- straightforward to include pQCD corrections

CP violation






- current bounds

$$\begin{aligned}d_e &< 8.7 \cdot 10^{-16} \text{ e fm} \\d_n &< 3.0 \cdot 10^{-13} \text{ e fm} \\d_{199\text{Hg}} &< 6.2 \cdot 10^{-17} \text{ e fm} \\d_{225\text{Ra}} &< 4.2 \cdot 10^{-17} \text{ e fm}\end{aligned}$$

- future bounds

$$\begin{aligned}d_e &< 5.0 \cdot 10^{-17} \text{ e fm} \\d_n &< 1.0 \cdot 10^{-15} \text{ e fm} \\d_{199\text{Hg}} &< 6.2 \cdot 10^{-17} \text{ e fm} \\d_{225\text{Ra}} &< 1.0 \cdot 10^{-14} \text{ e fm}\end{aligned}$$

Left-right symmetric model




- model based on $SU(3)_c \times SU(2)_L \times SU(2)_R \times U(1)_{B-L}$
- broken to SM group at $\nu_R \gtrsim 10 \text{ TeV}$
- generate ν masses via type-I and type-II see-saw

$K-\bar{K}$ oscillations and di-jet searches

need small Yukawas

Left-right symmetric model



- model based on $SU(3)_c \times SU(2)_L \times SU(2)_R \times U(1)_{B-L}$
- broken to SM group at $v_R \gtrsim 10 \text{ TeV}$


$K-\bar{K}$ oscillations and di-jet searches

- generate ν masses via type-I and type-II see-saw

need small Yukawas

- also generate dim. 7, with one Yukawa
- and dim. 9, with no Yukawa suppression

Left-right symmetric model



- model based on $SU(3)_c \times SU(2)_L \times SU(2)_R \times U(1)_{B-L}$
- broken to SM group at $\nu_R \gtrsim 10 \text{ TeV}$

$K-\bar{K}$ oscillations and di-jet searches

- generate ν masses via type-I and type-II see-saw

need small Yukawas

- also generate dim. 7, with one Yukawa
- and dim. 9, with no Yukawa suppression

Relation to charge-independence breaking




TABLE IX. Evolution of 1S_0 pp phase shifts from the charge-independent potential to the full interaction, as described in the text. Energies are in MeV.

T_{lab}	CI	+ m_p	+ CD v^*	+ CD v^R	+ v^{EM}
1	57.99	57.80	57.42	55.50	32.68
5	61.22	61.12	60.88	59.78	54.74
10	57.98	57.90	57.71	56.84	55.09
25	49.22	49.17	49.05	48.36	48.51
50	38.87	38.84	38.76	38.13	38.78
100	24.87	24.85	24.80	24.19	25.01
150	14.83	14.81	14.77	14.16	15.00
200	6.82	6.80	6.77	6.15	6.99
250	0.08	0.06	0.04	-0.60	0.23
300	-5.78	-5.79	-5.82	-6.47	-5.64
350	-10.99	-11.00	-11.01	-11.69	-10.86

AV18 potential, Phys. Rev. C51 (1995) 38-51

2. in realistic potentials (AV18, χ -EFT)

V_{CIB} and V_{CIB}^S give effects of comparable size (when Coulomb is perturbative)

- e.g. large $C_1 + C_2$ in χ -EFT potentials


$$\frac{C_1 + C_2}{2} \sim \frac{50}{(4\pi F_\pi)^2}$$

M. Piarulli et al, '16

- same effect in isotensor energy coeff. of light nuclei

Low-energy Effective Lagrangian for $\Delta L = 2$

$\Delta L = 2$ Lagrangian at 1 GeV




$$\mathcal{L}_{\Delta L=2} = \mathcal{L}_{\Delta L=2}^{\Delta e=0} + \mathcal{L}_{\Delta L=2}^{\Delta e=1} + \mathcal{L}_{\Delta L=2}^{\Delta e=2}$$

- $\mathcal{L}_{\Delta L=2}^{\Delta e=0}$ includes ν masses, magnetic moments, ...

$$\mathcal{L}_{\Delta L=2} = -\frac{1}{2}(m_\nu)_{ij} \nu_L^{Tj} C \nu_L^i + \dots \quad m_\nu \sim \mathcal{O}\left(\frac{v^2}{\Lambda}\right)$$

$\Delta L = 2$ Lagrangian at 1 GeV




$$\mathcal{L}_{\Delta L=2} = \mathcal{L}_{\Delta L=2}^{\Delta e=0} + \mathcal{L}_{\Delta L=2}^{\Delta e=1} + \mathcal{L}_{\Delta L=2}^{\Delta e=2}$$

- $\mathcal{L}_{\Delta L=2}^{\Delta e=1}$ starts at dim. 6, $C_i^{(6)} = \mathcal{O}\left(\frac{v^3}{\Lambda^3}\right)$

$$\mathcal{L}_{\Delta L=2}^{(6)} = \frac{2G_F}{\sqrt{2}} \left\{ C_{\text{VL}}^{(6)} \bar{d}_L \gamma^\mu u_L \nu_L^T C \gamma_\mu e_R + C_{\text{VR}}^{(6)} \bar{d}_R \gamma^\mu u_R \nu_L^T C \gamma_\mu e_R \right. \\ \left. + C_{\text{SL}}^{(6)} \bar{d}_R u_L \nu_L^T C e_L + C_{\text{SR}}^{(6)} \bar{d}_L u_R \nu_L^T C e_L + C_{\text{T}}^{(6)} \bar{d}_R \sigma^{\mu\nu} u_L \nu_L^T C \sigma_{\mu\nu} e_L \right\}$$

β decay w. the “wrong” neutrino & all possible Lorentz structures

$\Delta L = 2$ Lagrangian at 1 GeV



- $\mathcal{L}_{\Delta L=2}^{\Delta e=2}$ starts at dim. 9


$$\mathcal{L}_{\Delta L=2}^{(9)} = \frac{2G_F^2}{v} \left[\sum_{i=\text{scalar}} \left(C_i^{(9)} \bar{e}_L C \bar{e}_L^T + C_i^{(9)'} \bar{e}_R C \bar{e}_R^T \right) O_i + \bar{e}_R \gamma_\mu C \bar{e}_L^T \sum_{i=\text{vector}} C_{iV}^{(9)} O_i^\mu \right]$$

- a small set receives contributions from dim. 7 operators

$$C_1^{(9)}, C_{4,5}^{(9)} \sim \mathcal{O} \left(\frac{v^3}{\Lambda^3} \right)$$

- the remaining operators $\sim \mathcal{O} \left(\frac{v^5}{\Lambda^5} \right)$


Higher partial waves



- Weinberg's counting lead to problems in ${}^3P_{0,2}$ waves

\implies need LO counterterms in the strong interaction

Higher partial waves



- Weinberg's counting lead to problems in $^3P_{0,2}$ waves
 \implies need LO counterterms in the strong interaction
- neutrino potential in P waves does not require further renormalization